

Comprehensive Bibliography for the Pine Hill Area and Rare Plants

- Ayres, D. R. 1997. The clonal and population biology of *Wyethia reticulata* (Asteraceae), a rare perennial of the chaparral. Ph.D. dissertation, University of California, Davis, CA.
- Ayres, D.R. and Ryan, F.J. 1999. Genetic diversity and structure of the narrow endemic, *Wyethia reticulata*, and its congener *W. bolanderi* (Asteraceae) using RAPD and allozyme techniques. *American Journal of Botany* 86: 344-353
- Ayres, D.R., and Ryan, F.J. 1997. The clonal and population structure of a rare endemic plant, *Wyethia reticulata* (Asteraceae): allozyme and RAPD analysis. *Molecular Ecology* 6:761-772.
- Ayres, D.R. 2011. Effects of fire on the demography of three rare chaparral plants: (*Calystegia stebbinsii*, *Ceanothus roderickii*, and *Wyethia reticulata*). Proceedings of the CNPS Conservation Conference, 17–19 Jan 2009 pp. 1–9, California Native Plant Society
- Baad, M. F. and G. D. Hanna. 1987. Pine Hill Ecological Preserve operations and maintenance schedule. Prepared for California Depart of Fish and Game. Unpublished report.
- Barnes, J. J. 2001. Cultural resources reconnaissance of Bureau of Land Management lands along the South Fork along of the American River, El Dorado County, California. Report on file, Bureau of Land Management, Folsom Field Office, Folsom, California.
- Boyd, R. S. 2007. Response to fire of *Ceanothus roderickii* (Rhamnaceae), a federally endangered California endemic shrub. *Madroño* 54:13-21.
- Boyd, R. S. 2003. Factors affecting seed production by the endangered chaparral shrub *Fremontodendron californicum* ssp. *decumbens* (Sterculiaceae) *Madroño* 50: 232-242.
- Boyd, R. S. 2001. Ecological benefits of myrmecochory for the endangered chaparral shrub *Fremontodendron decumbens* (Sterculiaceae). *Am. J. of Botany* 88: 234-241.
- Boyd, R. S. 1996. Ant-mediated dispersal of the rare chaparral shrub *Fremontodendron decumbens* (Sterculiaceae). *Madroño* 43: 299-315.
- Boyd, R. S. 1994. Pollination biology of the rare shrub *Fremontodendron decumbens* (Sterculiaceae). *Madroño* 41: 277-289.
- Boyd, R. S. and L. L. Serafini. 1992. Reproductive attrition in the rare chaparral shrub *Fremontodendron decumbens* Lloyd (Sterculiaceae). *Am. J. of Botany* 89: 1264-1272.
- Brummitt, R.K. 1974. A remarkable new species of *Calystegia* (Convolvulaceae) from California. *Kew Bulletin* 29: 499-502.
- Brummitt, R.K and S. Namoff. 2013. *Calystegia vanzuukiae* (Convolvulaceae), a remarkable new species from central California. *Aliso* 31: 15-18

- Bureau of Land Management. 2008. Sierra Resource Management Plan and Record of Decision. Folsom Field Office, Folsom, California.
- Bureau of Land Management. 2008. Pine Hill Preserve Resource Management Plan. Folsom Field Office, Folsom, California.
- Bureau of Land Management. 2002. A brief history of the Pine Hill Preserve in El Dorado County, California. Folsom Field Office, Folsom, California. Unpublished.
- Burge , D. O. , D. M. Erwin , M. B. Islam , J. Kellermann , S. W. Kembel, Manos, P.S., and D. H. W. Wilken. 2011. Diversification of the genus *Ceanothus* (Rhamnaceae) in the California Floristic Province. *International Journal of Plant Sciences* 172: 1137 – 1164
- Burge , D. O., and P. S. Manos. 2011. Edaphic ecology and genetics of the gabbro-endemic shrub *Ceanothus roderickii* (Rhamnaceae). *Madrono* 58 : 1 – 21
- Burge et al 2012. Microsatellite markers from *Ceanothus roderickii* (Rhamnaceae) using Next-Generation sequencing technology. *American Journal of Botany*: e127–e130
- Burge et al 2013. Limited hybridization across an edaphic disjunction between the gabbro-endemic shrub *Ceanothus Roderickii* (Rhamnaceae) and the soil-generalist *Ceanothus cuneatus*. *American Journal of Botany* 100(9): 1883–1895.
- California Department of Fish and Game. 1999. Draft Management Plan for Pine Ecological Reserves, El Dorado County.
- EIP Associates. 1991. Preserve sites and preservation strategies for rare plan species in western El Dorado County. Unpublished.
- Gogol-Prokurat, M. 2011. Predicting habitat suitability for rare plants at local spatial scales using a species distribution model. *Ecological Applications* 21:33–47.
- Gogol-Prokurat, M. 2014. Characterizing habitat suitability for disturbance-dependent rare plants of gabbro soils. *California Fish and Game* 100: 19-33
- Greene E.L. 1883. New western Compositae. 1883. *Bulletin of the Torrey Botanical Club* 10: 86-89
- Greene, E.L. 1884. *Wyethia reticulata*. *Bull. Cal. Acad. Science* page 9.
- Hardig, T. M., P. S. Soltis and D. E. Soltis. 2000. Diversification of the North American shrub genus *Ceanothus* (Rhamnaceae): conflicting phylogenies from nuclear ribosomal DNA and chloroplast DNA. *Am. J. of Botany*. 87: 108-123.
- Horenstein, J.E. and A. Ehrgott. 1997. Pine Hill Ecological Reserve. A grant application submitted by the California State Department of Fish and Game and the American River Conservancy to the Bureau of Reclamation and the U.S. Fish and Wildlife Service under the Central Valley Project Improvement Act of October, 1992 (CVPIA, B-1 Other Program).

- James, S. C. 1996. A demographic study of *Ceanothus roderickii* (the Pine Hill Ceanothus) El Dorado County, California. M.Sc. thesis, California State University, Sacramento, CA.
- Kelman, W., L. Broadhurst, C. Brubaker and A. Franklin. 2006. Genetic relationship among *Fremontodendron* (Sterculiaceae) populations of the Central Sierra Nevada foothills of California. *Madroño* 53: 380-387.
- Lloyd, R. M. 1965. A new species of *Fremontodendron* (Sterculiaceae) from the Sierra Nevada Foothills, California. *Brittonia* 17: 3382-384.
- Marsh, G. 2000. Genetic structure of Layne's Butterweed (*Senecio layneae*) using random amplified polymorphic DNA (RAPD) and inter simple sequence repeat (ISSR) markers. MSc. thesis, California State University, Sacramento, CA.
- Marsh, G. D. and D. R. Ayres. 2002. Genetic structure of *Senecio layneae* (Compositae): A rare plant of the chaparral. *Madroño* 49:150-157.
- Medeiros I.D, Rajakaruna N., and Alexander E. 2015. Gabbro Soil-Plant Relations in the California Floristic Province. *Madroño*, 62(2):75-87.
- Nosal, T. R. D. 1997. A demographic study of Stebbins' Morning Glory, (*Calystegia stebbinisii*, Brummit, Convolvulaceae), a California state listed and federally listed endangered plant species. MSc. thesis, California State University, Sacramento, CA.
- Safford, H. D., J. H. Viers, and S. P. Harrison. 2005. Serpentine endemism in the California Flora: a database of serpentine affinity. *Madroño* 52:222-257.
http://www.des.ucdavis.edu/faculty/Harrison/people/sue/papers/madrono_paper.pdf
- United States Fish and Wildlife Service. 2004. A Draft Recovery Plan for Fifteen Plants of the Southern Sierra Nevada Foothills, California.
- United States Fish and Wildlife Service. 2002. Recovery Plan for Gabbro Soil Plants of the Central Sierra Nevada Foothills, California.
- United States Fish and Wildlife Service. 1996. Endangered and Threatened Wildlife and Plants; Determination of Endangered Status for Four Plants and Threatened Status for One Plant from the Central Sierra Foothills of California. *Federal Register* 61: 54346-54358.
- Weber, W.A. 1946. A Taxonomic and Cytological Study of the Genus *Wyethia*, Family Compositae, with Notes on the Related Genus *Balsamorhiza*. *American Midland Naturalist* 35: 400-452
- Wilson, J. L. 1986. A study of plant species diversity and vegetation patterns associated with the Pine hill gabbro formation and adjacent substrata, El Dorado County, California. M.Sc. thesis, California State University, Sacramento, CA.

Wilson, J. L., Ayres D. R., Steinmaus, S., and M. Baad. 2009. Vegetation and flora of a biodiversity hotspot: Pine Hill, El Dorado County, California, USA. *Madroño* 56:246–278.

Zika, P.F., Janeway, L.P., and B.L. Wilson. 2014. *Carex xerophila* (Cyperaceae), a new sedge from the chaparral of northern California. *Madroño*, 61(3):299-307